

PATIENT ALLERGIC TO:

DRAFT COPY

DOCTOR'S ORDER SHEET

(Submit approved copy to I.S.)

UNAPPROVED ABBREVIATIONSDO NOT USE THE FOLLOWING:					
			se of an unnecessary trailing zero		The use of a leading decimal point
		The u	se of an abbreviation for Morphine or Magnesium S	Sulfate	QOD qod IU U QD qd
DATE	HR	Cellulitis (approved June 2018)			
		1	Laboratory		
			□ C-Reactive Protein □ Now	□ In AM	
			□ CBC w/Diff □ Now	□ In AM	
			□ ESR □ Now	□ In AM	
		2.	Cultures		
		Blood x2 Wound Site:			
		3. Consults			
			General Surgeon, routine		Orthopaedic, routine
			Wound Care, routine		Podiatry, routine
		5.	Probiotics		
	IDSA Guidelines state there is insufficient data at this time to recommend probiotics for primary				
			prevention of CDI. Studies suggest they		
			Saccharomyces Boulardii 250m	÷	
		4.	Antibiotics (pharmacy will adjust do	,	(world icon link to one page tx guideline?)
				re B-lactam allergy use v	ancomycin)
			Cefazolin 1g IVPB q6h		
			Purulent Cellulitis (or MRSA Risk Vancomycin 1g IVPB x1	Factors)	
			 Pharmacist Consult- Vancomyci 	in	
		Cellulitis associated with sepsis, diabetic foot infections, necrosis or gangrene			
		(For severe B-lactam allergy, ok to use Meropenem)			
		□ Piperacillin/Tazobactam 3.375g IVPB Q8H			
			 Meropenem 1g IVPB Q8H 		
			□ Vancomycin 1g IVPB x1		
			Pharmacist Consult- Vancomyc	in	
		Oral Stepdown			
		Cephalexin 500 mg PO QID			
		Augmentin 875mg PO BID			
		Doxycycline 100 mg PO BID			
			Bactrim DS 1 tablet PO BID		
			□ Clindamycin 300 mg PO TID		
MD Signature				Tim	e / Date
UPDATED:				ORDER #:	
REVIE	NED				

DOCTOR'S ORDER